

①



Europäisches Patentamt
European Patent Office
Office européen des brevets

②

Publication number:

0 273 115
A2

③

EUROPEAN PATENT APPLICATION

④

Application number: 87114490.3

⑤

Int. Cl.⁴ C07D 219/16, C07D 221/12,
G01N 33/53, C09K 11/06

⑥

Date of filing: 05.10.87

The title of the invention has been amended
(Guidelines for Examination in the EPO, A-III,
7.3).

⑦

Applicant: ABBOTT LABORATORIES

Abbott Park Illinois 60064(US)

⑧

Inventor: Mattingly, Phillip Gregory
204 Seafarer Drive
Grayslake Illinois 60030(US)
Inventor: Bennett, Larry Gene
84 W. Belvidere Road
Grayslake Illinois 60030(US)

⑨

Priority: 22.10.86 US 921979

⑩

Date of publication of application:
06.07.88 Bulletin 88/27

⑪

Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

⑫

Representative: Modiano, Guido et al
MODIANO, JOSIF, PISANTY & STAUB
Modiano & Associati Via Meravigli, 16
I-20123 Milan(IT)


⑬

Chemiluminescent acridinium and phenanthridinium salts.

⑭

Acridinium sulfonylamides and isomers, such as phenanthridinium sulfonylamides, may be employed in applications including chemiluminescent immunoassays. Methods for synthesis of these compounds include contacting an amine with a sulfonylhalide to form a sulfonamide and acylating with an activated carboxylic acid of an acridine or isomer thereof. The N-sulfonyl-9-acridinium carboxamide and isomer may be conjugated to antigens, haptens, antibodies, and nucleic acids for use in chemiluminescent assays.

EP 0 273 115 A2



CHEMILUMINESCENT ACRIDINIUM SALTS

Background

The present invention relates in general to chemiluminescent methods and materials and in particular to methods and materials involving chemiluminescent acridinium and phenanthridinium salts.

Chemiluminescence may be defined as the generation of light from a chemical reaction. The mechanism of most chemiluminescent reactions is not known in detail, but a generalized mechanism [Schuster et al., Advances in Physical Organic Chemistry, 187-238 (1984)] may be outlined:



Compound A undergoes a chemical reaction (usually oxidation) to yield a product in an electronically excited State ("B^{*}"). As it returns to the ground state ("B"), this product gives up energy in the form of light ("hν").

Although competing dark reactions may decrease the efficiency of the overall reaction to less than 1%, some bioluminescent systems may achieve 60-70% efficiency, and, in many cases, limits of detection in the femtomole (10⁻¹⁵ mole) to attomole (10⁻¹⁸ mole) range have been recorded.

Chemiluminescence has been used for a variety of purposes in analytical chemistry where other methods fail to have adequate sensitivity. In immunodiagnostics, chemiluminescent immunoassays ("CLIA") may thus match or exceed the sensitivity of radioimmunoassays ("RIA") or enzyme immunoassays ("EIA") [Kircka et al., Diagnostic Medicine, 1, 45-52 (1984)].

Luminol and isoluminol derivatives are the most widely used chemiluminescent reagents for immunoassays. The light-yielding reaction is initiated by oxidation with alkaline hydrogen peroxide in the presence of catalysts such as microperoxidase or transition metal ions. Light emission occurs at about 465 nm, which corresponds to the fluorescence emission of the product, aminophthalic acid. Aminobutylethyl isoluminol ("ABEI") may be used as a label in immunoassays and is commercially available.

A second group of chemiluminescent reagents, aryl oxalates [Gill, Aldrichimica Acta, 16, 59-61 (1983) and Catherall et al., J. Chem. Soc. Faraday Trans. 2, 80, 823-834 (1984)], have been used as commercial cold light sources [see e.g., Tseng et al., U.S. Patent No. 4,338,213] and in high performance liquid chromatography ("HPLC") detectors [Kobayashi et al., Anal. Chem., 52, 424-427 (1980) and Miyaguchi et al., J. Chromatogr., 303, 173-176 (1984)]. It is thought that these derivatives react with hydrogen peroxide in buffered or unbuffered solvents to give a dioxetan-dione which decomposes quickly to give CO₂ in an excited state. Energy is then transferred by electron transfer to a fluorescer molecule which emits light.

A third group of reagents, 10-methyl-acridinium-9-carboxylic acid aryl esters, are chemiluminescent in the presence of alkaline hydrogen peroxide and in the absence of a catalyst. The mechanism is thought to involve initial attack by a hydroperoxide anion, followed by intramolecular displacement of the phenolate (the "leaving group") to give a strained dioxetan-one. The strained dioxetan-one decomposes to CO₂ and excited N-methyl-acridone, which emits light at 430 nm. Carboxy-substituted acridinium salts have been used as labels in immunoassays [Weeks et al., Clin. Chem., 29, 1474-79 (1983); Campell et al., European Patent Application No. 82,636; and McCapra et al., UK Patent No. GB 1,461,877]. Also, 5-methyl-phenanthridinium-6-carboxylic acid aryl esters, which are isomeric with the acridinium aryl esters, have been used as labels in immunoassays [Lin et al., European Patent Application No. 170,415].

Despite their usefulness in immunoassays, antibody-conjugated phenyl 10-methyl-9-acridiniumcarboxalates, in our hands, are unstable due to hydrolysis above pH 4.0 (-20°C to 40°C), losing greater than 10% of their activity within three days. Although acridinium esters are stable below pH 4.0, conjugate antibodies are often not stable in this pH range.


In Tseng et al., supra, bis-N-alkyl-N-trifluoromethyl sulfonyl oxalamides are indicated to be more stable than the corresponding aryl esters and are also indicated to be as efficient. The nucleofugacity of the phenol and the trifluoromethyl sulfonamide are indicated to be comparable, i.e. it is indicated that each has a pK_a of about 7. Gill, supra, "look forward" to the development of a particular sulfonyl oxalamide as an example of an oxalate with "higher" quantum efficiency.

Brief Description of the Drawings


The Figure illustrates the synthesis of a 10-alkyl-N-sulfonyl-9-acridinium carboxamide according to the present invention.

Summary of the invention

The present invention provides chemiluminescent compounds identified by the formula



and isomers thereof including isomers identified by the formula



wherein R, R', R'', X¹, X², and X³ are substituents which do not interfere with effective chemiluminescence, with the proviso that R-X³, R'-X² and R''-X¹ may be independently hydrogen. More specifically, R, R' and R'' may be spacer arms and X¹, X² and X³ may be independently members of the group consisting of hydrogen, carboxy, carboalkoxyl, carboxamido, carboaryloxy, cyano, carboximido, isocyanato, isothiocyanato, sulfo, sulfonyl halide, carbonyl halide, N-succinimidyloxycarbonyl and N-maleimide groups. Y is an appropriate counterion and may be selected from the group consisting of sulfate, alkylsulfate, halosulfate, haloborate, haloacetate, halophosphate, phosphate and halide.

R, R', and R'' may independently include a member selected from the group consisting of alkyl, alkylene, aryl, substituted alkyl, substituted alkylene and substituted aryl groups, such that one or more hydrogens of said member is replaced by an alkyl, aryl, alkylene, substituted alkyl, substituted alkylene, substituted aryl, alkoxy, aryloxy, halo, amino, protected amino, substituted amino hydroxy, protected hydroxy, oxo, thio, imino, mercapto or substituted mercapto group; or such that one or more carbon atoms of the member is replaced by a heteroatom. The heteroatom may be selected from the group consisting of nitrogen, phosphorus, sulfur and oxygen.

R, R', and R'' independently may also be spacer arms of the formula $-(CH_2)_n-$


where $n = 0 - 50$. Specifically, R'' may be $-CH_2-$ and X¹ may be $-H$.

The currently most preferred compounds according to the present invention for use in chemiluminescent immunoassays are 10-methyl-N-[2-carboxyethyl]-N-tosyl-9-acridinium carboxamide, 10-(3-sulfopropyl)-N-(2-carboxyethyl)-N-tosyl-9-acridinium carboxamide and 1-(3-sulfopropyl)-N-(3-sulfopropyl)-N-tosyl-9-acridinium carboxamide.


A method, according to the present invention, for preparation of a chemiluminescent compound includes the steps of contacting an amine identified by the formula



with a sulfonylhalide identified by the formula



in an inert solvent in the presence of base to form a sulfonamide anion and metal ion identified by the formulas



and acrylating with an activated 9-acridinecarboxylate compound according to the present invention, wherein W is selected from the group: consisting of chloro and fluoro groups, wherein M is selected from the group consisting of Li, Na and K, wherein the activating group is selected from the group consisting of halo, imidazo, N-hydroxysuccinimidyl and azido groups and wherein all other symbols are as defined above.

A conjugate according to the present invention may be formed by covalently coupling an antibody, a hapten, an antigen or a polynucleotide (e.g., DNA or RNA) to a chemiluminescent compound according to the present invention, and a method for performing a chemiluminescent assay comprises the step of exposing a sample to be tested to the conjugate in order to detect the presence of a substance specifically reactive with the conjugate, e.g., a specific antigen, a specific antibody or a complementary polynucleotide (i.e., a polynucleotide which forms sequence-specific hydrogen bonds with the polynucleotide conjugate according to the present invention).

Detailed Description

The problem of acridinium aryl ester instability is approached in the present invention by changing the leaving group from a phenolate to a sulfonamide anion. While both leaving groups have a pK_a of about 10, the acridinium sulfonylamide has the additional stabilization associated with amide bonds. This is reflected in a comparison in the infrared of the carbonyl stretching frequency of the aryl ester (1730 cm^{-1}) with that of the sulfonylamide (1680 cm^{-1}).

A class of acridinium salts, 10-alkyl N-alkyl (aryl) sulfonyl-N-alkyl(aryl) 9-acridinium carboxamide salts, was prepared according to the general scheme illustrated in the Figure. In the Figure, R, R' and R'' are substituents which may function as spacer arms, solubility modifiers and/or reactivity modifiers but which do not interfere with the chemiluminescent reaction. ("Interfere" is defined herein to mean "prevent the production of effective chemiluminescence", i.e., prevent production of chemiluminescence to the extent that the compound is not useful for the intended application.) Also in the Figure, X', X², X³ are substituents which may function as solubility enhancers and/or as reactive groups for linkage to an analyte or as groups which may be readily converted to such reactive or linker groups by means well known to those skilled in the art. Y is a counterion in the Figure.


Salts produced according to the scheme of the Figure have generated light upon oxidation with alkaline hydrogen peroxide. The compounds were made from readily available amines ($\text{X}^3\text{-RNH}_2$) and sulfonyl chlorides ($\text{X}^2\text{-R'-SO}_2\text{Cl}$). When acrylated with 9-chlorocarbonyl acridine, the intermediate sulfonamide ($\text{X}^3\text{-RNH-SO}_2\text{R'-X}^2$) gave a new class of acridine compounds, which on alkylation gave the acridinium salts. Similarly, substitution of a 6-chlorocarbonyl phenanthridine for the acridine in this scheme gives rise to a new class of phenanthridinium salts. These acridinium and phenanthridinium salts are useful for chemiluminescent labeling of proteins, nucleic acids and small molecules used in diagnostic testing.

Several acridinium sulfonylamides were prepared which have specific activity and stability suitable for use in diagnostic testing, particularly in CLIA. The synthesis of these compounds allows for the introduction of a variety of functional groups (X', X², X³) which may be used in antibody labeling. In addition, the kinetics of the chemiluminescent reaction may be controlled by the choice of the substituents (R, R') on the sulfonamide leaving group.

The compounds were evaluated for their efficiency by diluting 20 μl of a 10^{-3} M solution of the compound with 300 μl of 0.1N HCL, then adding 150 μl of 0.03% H_2O_2 in 0.2 N NaOH to trigger the chemiluminescence. Chemiluminescence was measured on a photon-counting luminometer. The light output was recorded as total photon counts, from which the efficiency of each compound was calculated as counts/mole. These are relative numbers, since the efficiency of the photon counting was instrument-dependent. Direct comparisons of compounds were carried out on the same instrument. The results are presented in Table I which the structures may be identified by the formula

5

10



Y -

wherein $R''-X'$ is CH_3 , and $R'-X^2$ and $R-X^3$ are as indicated in Table 1, chemiluminescent output is abbreviated "CTS.MOLE." the time required for total light output is abbreviated "INT. TIME" and the time required to reach peak light output is abbreviated "PEAK CTS."

20

25

30

35

40

45

50

55

TABLE 1

CTS/MOLE

$\frac{R-X^3}{R'-X^2}$	$\frac{CTS}{MOLE}$ ($\times 10^{-18}$)	INT. TIME	PEAK CTS (SEC)
CF ₃	12	1	0.22
O-NO ₂ C ₆ H ₄	10	2	0.23
p-Br-C ₆ H ₄	9	2	0.24
CF ₃	15	2	0.25
p-NO ₂ -C ₆ H ₄	8	2	0.25
O-NO ₂ -C ₆ H ₄	11	2	0.25
p-CH ₃ C ₆ H ₄	9	2	0.27
O-NO ₂ C ₆ H ₄	6	2	0.29
2,4-di-NO ₂ C ₆ H ₃	5	2	0.32
p-BrC ₆ H ₄	7	3	0.44
p-BrC ₆ H ₄	12	6	0.44
p-CH ₃ C ₆ H ₄	5	6	0.98
p-CH ₃ C ₆ H ₄	8.3	10	0.96
2,4,6-(C ₃ H ₇) ₃ C ₆ H ₂	14	20	4.08
2,4,6-(CH ₃) ₃ C ₆ H ₂	5	50	11.6
CF ₃	4	2	--

All of the tested compounds were efficient ($5-20 \times 10^4$ counts/mole). The specific activity was insensitive to the nature of the R and R' groups at locations indicated above; however, the time required to reach peak light output and the time required for total light output differed by a factor of 50 between the fastest and slowest compounds. Electron withdrawing groups in R and R' increased the reaction rate while bulky, electron-donating groups decreased the reaction rate. Although chemiluminescent compounds according to the present invention which have a chemiluminescent lifetime of 2-10 seconds are preferred for immunoassays, compounds having shorter lifetime may be useful as a source of intense, pulsed light, and compounds having a longer lifetime may be useful as "cold light" sources.

The stability of compounds prepared according to the present invention was assessed in several ways. First, the compounds were diluted to sub-nanomolar solutions in aqueous buffer at pH 5-7. The solutions

U 273 115
were incubated at room temperature and at 45°C, while the decrease in chemiluminescence was monitored over time. This provided qualitative results whereby the relative stability of the compounds was determined. Anomalous results due to non-specific adsorption of the compounds on the incubation container were minimized by the addition of detergents, protein, and the like. Unambiguous, quantitative results were obtained by monitoring millimolar solutions of the compounds by reverse phase high performance liquid chromatography ("HPLC"). The stability of these compounds was affected by R and R' in the same way as were the kinetics of the chemiluminescence reaction, i.e. electron withdrawing groups destabilized and bulky electron donating groups stabilized the compounds.

Although other techniques may be employed to label antibodies, the NHS activation method is presently preferred. Other materials which function well according to the present invention include polyclonal antibodies, monoclonal antibodies, Fab antibody fragments, all of which are hereinafter included in the general term "antibody," haptens, antigens, nucleic acid probes, and non-antibody binding proteins capable of binding complementary small molecular weight analytes (for example, folate binding protein, which binds folic acid, and intrinsic factor, which binds Vitamin B₁₂). Antibody conjugates retain more than 80% chemiluminescence after being heated at 45°C for four weeks.

A solid phase sandwich immunoassay system for assaying hepatitis B surface antigen ("HBsAg") (Abbott Laboratories, Abbott Park, Illinois) was employed to compare CLIA according to the present invention with RIA. The type of antibody-coated bead, diluent, incubation conditions, washing condition and antibody preparation were the same except that the antibody was labeled with ¹²⁵I by the chloramin T method for RIA and labeled with NHS-activated N-sulfonyl-9-acridinium carboxamide for CLIA.

A solid phase sandwich immunoassay for human thyroid stimulating hormone (hTSH) was used to compare CLIA with EI (Abbott Laboratories, Abbott Park, Illinois). The EIA employed a horseradish peroxidase ("HRPO")-labelled antibody while the CLIA used an NHS-activated N-sulfonyl-9-acridinium carboxamide.

The present invention is more specifically described in the following examples. In Example 1, the preparation of sulfonamides which are useful in constructing compounds according to the present invention is set forth. Example 2 includes a description of the preparation of N-sulfonyl-9-acridinecarboxamides according to the present invention. In Example 3, the preparation of 10-methyl N-sulfonyl-acridinium carboxamides is described. Examples 4-6 contain descriptions of syntheses of p-toluenesulfonyl (tosyl) compounds according to the present invention. In Example 7, the preparation of acridinecarboxamides is illustrated.

Example 8-10 contain methods for synthesis of some acridinium carboxamides and products thereof according to the present invention. In Example 11, an evaluation of the chemiluminescence of N-sulfonylacridinium carboxamide compounds according to the present invention is provided. Example 12 includes a report of a stability test of an acridinium carboxamide according to the present invention. In Example 13, the temperature and pH stability of two acridinium carboxamides according to the present invention is compared to the temperature and pH stability of an acridiniumcarboxylate. Example 14 is a description of a method for conjugating an antibody, specifically an immunoglobulin G ("IgG") antibody, with a compound according to the present invention. The results of a heat stability study of a conjugate according to Example 14 are presented in Example 15. Example 16 includes a description of the preparation of anti-HBsAg acridinium-labeled conjugate as well as a comparison of the sensitivity observed in CLIA and RIA assays employing those conjugates. In Example 18, the synthesis of a phenanthridinium compound according to the present invention is described. Example 17 describes an anti-hTSH acridinium-labeled conjugate along with a comparison to an EIA system.

Example 1

General Method for Preparation Of Sulfonamides

Amine starting materials for compounds 1-13 and 17-21 are available from Aldrich Chemical Co., Milwaukee, Wisconsin. For compounds 14-16 and 22-25, the appropriate aminocarboxylic acid (as obtained from Aldrich Chemical Co., Milwaukee, Wisconsin) was esterified according to standard, published procedures to provide the starting materials.

In order to prepare a sulfonamide according to the present invention, the corresponding amine (200 mole percent) was dissolved in anhydrous methylene chloride, and was treated dropwise at 0°C with a solution (100 mole percent) of the sulfonyl chloride or anhydride. The solution was poured into anhydrous ether (5 volumes), washed with 1.4 M H₃PO₄ (25 ml) and then brine (25 ml), and dried over MgSO₄. After

filtering and evaporating, crude sulfonamides were crystallized from an appropriate solvent.

The following sulfonamides were prepared in this manner. In the description accompanying the name of each compound, the abbreviation "MS" identifies peaks, such as the base peak ("M⁺") in the mass spectrum at a location (i.e., at an m/e) specified by the symbol "@". A melting point ("M_p") or an indication that the material is a liquid at room temperature (e.g. "oil") or decomposes before melting ("decomp.") may be provided. Each compound is identified by a "compound number" (1-25 in this Example) followed by an "identifying number" (e.g. 13513-227) and a chemical name.

10

15

20

25

1. 13513-227 N-Phenyl-p-toluenesulfonamide
MS M⁺ @ 247
M_p 100-102°C

30

2. 13513-228 N-Phenyl-p-bromobenzenesulfonamide
MS M⁺ @ 311
M_p 115-117°C

35

40

45

50

55

3. 13513-229 N-Phenyl-o-nitrobenzenesulfonamide
MS M^+ @ 278
 M_p 112-113°C
4. 13513-231 N-Phenyl-p-nitrobenzenesulfonamide
MS M^+ @ 278
 M_p 168-170°C
5. 13513-232 N-Phenyl-2,4-dinitrobenzene-
sulfonamide
MS M^+ @ 323
 M_p 110-113°C
6. 13513-233 N-Phenyl-trifluoromethane-
sulfonamide
MS M^+ @ 225
 M_p 65-67°C
7. 13514-001 N-Isopropyl-p-
toluenesulfonamide.
MS M^+ @ 213
 M_p 50-51°C
8. 13514-002 N-Isopropyl-p-
bromobenzenesulfonamide
MS M^+ @ 277
 M_p 95-96°C
9. 13514-003 N-Isopropyl-o-
nitrobenzenesulfonamide
MS M^+ @ 244
 M_p 119-120°C

- 5
10
15
20
25
30
35
40
45
50
55
10. 13514-004 N-Isopropyl-
trifluoromethanesulfonamide
MS (M - 1) @ 190
oil
11. 13514-006 N-Isopropyl-p-
nitrobenzenesulfonamide
MS M⁺ @ 244
M_p 113-114°C
12. 13514-025 N-Butyl-2,4,6-
trimethylbenzenesulfonamide
MS M⁺ @ 255
M_p 45°C
13. 13514-026 N-Butyl-2,4,6,-
trisopropylbenzenesulfonamide
MS M⁺ @ 339
M_p 104°C
14. 13514-032 Benzyl 6-(N-tosylamino)-
hexanoate
MS M⁺ @ 375
oil
15. 13514-057 t-Butyl N-tosyl-s-alanine
MS M⁺ @ 242 (M - 57)
oil
16. 13514-058 Benzyl 5-(N-tosylamino)-pentanoate
MS M⁺ @ 361
oil
17. 13513-170 N-Butyl-p-toluenesulfonamide,
MS M⁺ @ 227
M_p 42-44°C

- 5 18. 13513-173 N-Butyl-p-bromobenzenesulfonamide,
 MS M^+ @ 241
 M_p 53-54°C
- 10 19. 13513-172 N-Butyl-o-nitrobenzenesulfonamide,
 MS M^+ @ 258
 M_p 58-60°C
- 15 20. 13513-174 N-Butyl-p-nitrobenzenesulfonamide
 MS M^+ @ 258
20 M_p 80-81°C
21. 13513-213 N-Butyl-2,4-dinitrobenzene
 sulfonamide,
25 MS M^+ @ 304
 M_p 60-62°C
- 30 22. 13513-085 Benzyl 6-(N-trifluoromethyl-
 sulfonylamino)-hexanoate
 oil
- 35 23. 13513-083 Benzyl N-(trifluoromethylsulfonyl)-
 4-(carboxymethyl) aniline
- 40 24. 14973-1A Benzyl N-(5-carboxypentyl)-p-
 bromobenzenesulfonamide
45 MS M^+ @ 439
 M_p 52-56°C
- 50 25. 14973-37A Benzyl N-(5-carboxypentyl)-p-
 nitrobenzenesulfonamide
 MS M^+ @ 406
55 M_p 86-88°C

Exempl 2

Preparation of N-sulfonyl-9-acridinecarboxamides

Freshly sublimed potassium tert-butoxide (200 mole percent) and tri-n-butylbenzylammonium bromide (1 mole percent) were suspended in toluene under nitrogen. A selected sulfonamide (200 mole percent) was added, the mixture was stirred for 10-30 minutes before evaporating to dryness and the dried material resuspended in the solvent. [Alternatively, the phase transfer catalyst may be omitted and an appropriate anion may be generated in tetrahydrofuran.] After the addition of 9-chlorocarbonylacridine hydrochloride (200 mole percent), the reaction mixture was stirred for 3 to 14 hours at room temperature until no further change was noted by thin-layer chromatography ("TLC"). The reaction solution was diluted with ethyl ether (10 volumes) and washed with brine (25 ml). After drying over $MgSO_4$, filtering and evaporating, the crude product was chromatographed (on a Chromatotron™ chromatograph [available from Harrison Research, Palo Alto, California] using a 2 mm silica rotor and employing an ethylacetate/hexane gradient). The fractions containing the product were collected, evaporated and crystallized from ether/heptane (i.e., the fractions were dissolved in ether followed by the addition of heptane until the mixture became cloudy).

The following compounds were prepared from starting materials as indicated in brackets wherein starting materials prepared herein are identified by the number associated with them in Example 1 or in this example, and wherein a commercial source is provided in brackets for each identified starting material not synthesized herein. All other notations are explained in Example 1.

20

25

30

35

40

45

50

55

5	26. 13513-234	N-Phenyl-N-p-toluenesulfonyl- 9-acridinecarboxamide [compound 1] MS M^+ @ 452 M _p 200°C
10		
15	27. 13513-236	N-Phenyl-N-p-bromobenzene- sulfonyl 9-acridinecarboxamide [compound 2] MS M^+ @ 516 M _p 218-219°C
20		
25	28. 13513-240	N-Phenyl-N-o-nitrobenzene- sulfonyl 9-acridinecarboxamide [compound 3] MS M^+ @ 483 M _p 197-200°C
30		
35	29. 13513-242	N-Phenyl-N-p-nitrobenzene- sulfonyl-9-acridinecarboxamide [compound 4] MS M^+ @ 483
40		
45	30. 13513-243	N-Phenyl-N-trifluoromethane- sulfonyl-9-acridinecarboxamide [compound 6] MS M^+ @ 430 M _p 162°C
50		
55	31. 13514-007	N-Isopropyl-N-p-toluene- sulfonyl-9-acridinecarboxamide [compound 7] MS M^+ @ 418 M _p 163-164°C

5 32. 13514-009 N-Isopropyl-N-p-
 bromobenzenesulfonyl-9-
 acridinecarboxamide
 [compound 8]
10 MS M⁺ @ 482
 M_p 205°C

15 33. 13514-012 N-Isopropyl-N-o-nitrobenzene-
 sulfonyl-9-acridinecarboxamide
 [compound 9]
 MS M⁺ @ 449
20 M_p 215°C

25 34. 13514-001 N-Isopropyl-N-trifluoromethane
 sulfonyl-9-acridinecarboxamide
 [compound 10]
 MS M⁺ @ 396

30 35. 13514-028 N-Butyl-N-2,4,6,-trimethyl-
 benzenesulfonyl-9-acridine-
 carboxamide
35 [compound 12]
 MS M⁺ @ 460
 M_p 88-90°C

40 36. 13514-031 N-Butyl-2,4,6-triisopropylbenzene-
 sulfonyl-9-acridinecarboxamide
 [compound 13]
45 MS M⁺ @ 544

50 37. 13514-042 Benzyl N-tosyl-N-(5-carboxypentyl)-9-
 acridinecarboxamide
 [compound 14]
 MS M⁺ @ 550
55 oil

38. 13514-062 Benzyl N-tosyl-N-(4-carboxybutyl)-9-
acridinecarboxamide
[compound 16]
MS M^+ @ 566

39. 13514-069 t-Butyl N-tosyl-N-(2-carboxyethyl)-
9-acridinecarboxamide
[compound 15]
MS M^+ 504
 M_p 157-158°C

40. 13513-186 N-Butyl-N-p-toluenesulfonyl-9-
acridinecarboxamide
[compound 17]
MS M^+ @ 432
 M_p 122-123°C

41. 13513-191 N-Butyl-N-o-nitrophenylsulfonyl
-9-acridinecarboxamide
[compound 19]
MS M^+ @ 463
 M_p 170°C

42. 13513-195 N-Butyl-N-p-nitrophenylsulfonyl-9-
acridinecarboxamide
[compound 20]
MS M^+ 463
 M_p 210°C

43. 13513-218 N-Butyl-N-(2,4-dinitrophenylsulfonyl)
-9-acridinecarboxamide
[compound 21]
MS M^+ @ 508
 M_p 95°C

44. 14973-9C

Benzyl N-(5-carboxypentyl)-N-p-
bromobenzenesulfonyl-9-
acridinecarboxamide

[compound 24]

MS (M + H) @ 645

45. 14973-40C

Benzyl N-(5-carboxypentyl)-N-p-
nitrobenzenesulfonyl-9-
acridinecarboxamide

[compound 25]

MS (M + H) @ 645

46. 14973-88A

N-p-Toluenesulfonyl-9-
acridinecarboxamide

[p-toluene sulfonamide (Aldrich)]

M_p 276°C

47. 14973-21C

N-Allyl-N-p-toluenesulfonyl-9-
acridinecarboxamide

[compound 46]

M_p 136-138°C

48. 13513-202

N-Butyl-N-p-bromobenzenesulfonyl-
9-acridinecarboxamide

MS M⁺ @ 496/498M_p 148-149°CExample 3

45 Preparation of 10-Methyl N-sulfonylacridinium carboxamides

Methylation of N-sulfonylacridine carboxamides was performed according to the following procedure. Each acridine sulfonylamide was dissolved in anhydrous methylene chloride. Anhydrous Na₂CO₃ (5 X weight of the sulfonamide) was added followed by methyl triflate (20 X weight of the sulfonamide). The suspension
50 was stirred under nitrogen for 14-48 hours at room temperature to 40°C. The reaction was monitored by TLC (reverse phase). The product was obtained after filtration and evaporation of the solvent and of excess methyl triflate. Purification was achieved by triturating the solid residue with hot benzene or by reverse phase HPLC.

The following compounds were prepared, and they are described according to the numerals, symbols
55 and abbreviations which are explained in Example 1 or in Example 2.

5

10

15

20

25

30

35

40

45

50

55

49. 13513-246 10-Methyl-N-phenyl-N-p-
toluenesulfonyl-9-
acridinium carboxamide
trifluoromethanesulfonate
[compound 26]
MS M^+ @ 467
 M_p 210-24°C (decomp.)

50. 13513-247 10-Methyl-N-phenyl-N-p-
bromobenzenesulfonyl-9-
acridinium carboxamide
trifluoromethanesulfonate
[compound 27]
MS M^+ @ 531, 533
 M_p 240°C (decomp.)

51. 13513-248 10-Methyl-N-phenyl-o-nitro-
benzenesulfonyl-9-
acridinium carboxamide
trifluoromethanesulfonate
[compound 28]
MS M^+ @ 490
 M_p 248-50°C (decomp.)

5

10

15

20

25

30

35

40

45

50

55

52. 13513-249 10-Methyl-N-phenyl-N-
trifluoromethanesulfonyl-9-
acridinium carboxamide
trifluoromethanesulfonate
[compound 30]
MS M^+ @ 445

53. 13513-250 10-Methyl-N-phenyl-p-
nitrobenzenesulfonyl-9-
acridinium carboxamide
trifluoromethanesulfonate
[compound 29]
MS M^+ @ 484

54. 13514-013 10-Methyl-N-isopropyl-N-p-
toluenesulfonyl-9-
acridinium carboxamide
trifluoromethanesulfonate
[compound 31]
MS M^+ @ 433
 M_p 214°C

55. 13514-014 10-Methyl-N-isopropyl-N-p-
bromobenzenesulfonyl-9-
acridinium carboxamide
trifluoromethanesulfonate
[compound 32]
MS M^+ @ 497/499
 M_p 200°C (decomp)

56. 13514-018 10-Methyl-N-isopropyl-N-o-nitrobenzenesulfonyl-9-acridinium carboxamide trifluoromethanesulfonate [compound 33] MS M^+ @ 464
57. 13514-021 10-Methyl-N-isopropyl-N-trifluoromethanesulfonyl-9-acridinium carboxamide trifluoromethanesulfonate [compound 34] MS M^+ @ 411
58. 13514-037 10-Methyl-N-butyl-N-(2,4,6-trimethylbenzenesulfonyl-9-acridinium carboxamide trifluoromethanesulfonate [compound 35] MS M^+ @ 475 M_p 227°C (decomp.)
59. 13514-038 10-Methyl-N-butyl-N-(2,4,6-triisopropylbenzenesulfonyl-9-acridinium carboxamide trifluoromethanesulfonate [compound 36] MS M^+ @ 559 M_p 231°C (decomp.)
60. 13514-044 Benzyl 10-methyl-N-tosyl-N-(5-carboxypentyl)-9-acridinium carboxamide trifluoromethanesulfonate [compound 37]

5

10

15

20

25

30

35

40

45

50

55

61. 13514-079 t-Butyl 10-methyl-N-tosyl-
N-(2-carboxyethyl)-9-
acridinium carboxamide
trifluoromethanesulfonate
[compound 39]
MS M^+ @ 519
 M_p 207°C (decomp.)

62. 13513-211 10-Methyl-N-butyl-N-p-
toluenesulfonyl-9-
acridinium carboxamide
trifluoromethanesulfonate.
[compound 40]
MS M^+ @ 447

63. 13513-212 10-Methyl-N-butyl-N-p-
bromobenzenesulfonyl-9-
acridinium carboxamide
trifluoromethanesulfonate
[compound 48]
MS M^+ @ 511
 M_p 126°C

64. 13513-215 10-Methyl-N-butyl-N-o-
nitrophenylsulfonyl-9-
acridinium carboxamide
trifluoromethanesulfonate
[compound 41]
MS M^+ @ 478
 M_p 232-234°C

5

10

15

20

25

30

35

40

45

50

55

65. 13513-216 10-Methyl-N-butyl-N-p-nitrophenysulfonyl-9-acridinium carboxamide trifluoromethanesulfonate [compound 42]
MS M^+ @ 478
 M_p 201°C
66. 13513-230 10-Methyl-N-butyl-N-(2-4 dinitrophenylsulfonyl)-9-acridinium carboxamide trifluoromethanesulfonate [compound 43]
MS M^+ @ 523
 M_p 215-220°C
67. 14973-31B 10-Methyl-N-allyl-N-p-toluenesulfonyl-9-acridinium carboxamide trifluoromethanesulfonate [compound 47]
MS $M + 2$ @ 433
68. 14973-47A Benzyl 10-methyl-N-(5-carboxypentyl)-N-p-nitrobenzenesulfonyl-9-acridinium carboxamide trifluoromethanesulfonate [compound 45]
MS M^+ @ 626
 M_p 139-141°C

69. 14973-90A 10-Methyl-N-methyl-N-p-
toluenesulfonyl-9-
acridinium carboxamide
trifluoromethanesulfonate

[compound 46]

MS M⁺ @ 405

70. 14973-25A Benzyl 10-methyl-N-(5-carboxypentyl)-N-
(o-bromobenzenesulfonyl)-9-
acridinium carboxamide

[compound 44]

Example 4

Synthesis of 10-methyl-N-tosyl-N-(6-hexanoyl-N-hydroxysuccinimido)-9-acridinium carboxamide trifluoromethanesulfonate

Compound 37 (450 mg, 0.78 mmoles) was treated with 6 ml of 31% HBr in acetic acid at 50°C for 2 hours under N₂. The solution was poured into 30 ml of water and cooled. Carboxylic acid compound 71, 13514-045 [N-tosyl-N-(5-carboxypentyl)-9-acridinecarboxamide] was separated by filtration.

Compound 71 (100 mg., 0.2 mmol) was dissolved in dry methylene chloride (5 ml) and treated with N-hydroxysuccinimide (23 mg, 0.2 mmol) and dicyclohexylcarbodiimide (41 mg) under N₂ for 12 hours. After reacting, the solution was filtered and then evaporated to dryness to yield an active ester, compound 72, 13514-952 [N-tosyl-N-(6-hexanoyl-N-hydroxysuccinimido)-9-acridinecarboxamide].

Compound 72 was methylated as in Example 3 to give compound 73. Compounds 71, 72 and 73 are described below using the numerals, symbols and abbreviations which are explained in Example 1.

explained in Example 1.

71. 13514-045 N-Tosyl-N-(5-carboxypentyl)-9-acridinecarboxamide

[compound 37]

MS M⁺ @ 240

M_p 150-152°C

72. 13514-052 N-Tosyl-N-(6-hexanoyl-N-hydroxysuccinimido)-9-acridinecarboxamide

[compound 71]

MS M⁺ @ 588

73. 13514-054 10-Methyl-N-tosyl-N-(6-hexanoyl-N-hydroxysuccinimido)-9-acridinumcarboxamide trifluoromethanesulfonate

[compound 72]

Example 5

Synthesis of 10-Methyl-N-tosyl-N-(5-pentanoyl-N-hydroxysuccinimido)-9-acridinium trifluoromethanesulfonate

Compound 38, 13514-062, was treated as in Example 4 and yielded compound 74, 13514-065 [N-tosyl-N(4-carboxybutyl)-9-acridinecarboxamide].

Compound 74 was coupled to N-hydroxysuccinimide, as in Example 4, to give compound 75, 13514-067, N-tosyl-N-(5-pentanoyl-N-hydroxysuccinimido)-9-acridinecarboxamide. This compound was methylated as in Example 3 to give compound 76, 13514-78 [10-methyl N-tosyl-N-(5-pentanoyl-N-hydroxysuccinimide)-9-acridinium carboxamide trifluoromethanesulfonate].

Compounds 74, 75 and 76 are described using the numerals, symbols and abbreviations which are explained in Example 1.

74. 13514-065 N-Tosyl-N-(4-carboxybutyl)-9-
 5 acridinecarboxamide
 MS M^+ @ 476
 M_p 152-155°C
75. 13514-067 N-Tosyl-(5-pentanoyl N-hydroxy
 10 succinimido)-9-acridinecarboxamide
 [compound 74]
 15 MS M^+ @ 573
76. 13514-078 10-Methyl-N-tosyl-N-(5-
 20 pentanoyl-N-hydroxy-
 succinimido)-9-
 acridinium carboxamide
 25 trifluoromethanesulfonate
 [compound 75]

Example 6

- 30 Synthesis of 10-methyl-N-tosyl-N-(2-carboxyethyl-9-acridinium carboxamide trifluoromethanesulfonate

Compound 61, 13514-079 (50 mg, 0.072 mmol) was dissolved in 2 ml of trifluoroacetic acid ["TFA"] at 0°C under N_2 . After stirring for 15 minutes, the TFA was evaporated and the residue was recrystallized from methanol/ether (i.e., the residue was dissolved in methanol, adding ether until cloudy). Alternatively,
 35 compound 61, was refluxed in 1 N HCl for 3 hours. The aqueous solution was evaporated to dryness to leave a residue, and the residue was purified by preparative reverse phase HPLC. Compound 77, 13514-081 [10-methyl N-tosyl-N-(2-carboxyethyl)-9-acridinium carboxamide] resulted from either approach. Compound 77 is described using the numerals, symbols and abbreviations which are explained in Example 1.

- 40 77. 13514-081 10-Methyl-N-tosyl-N-
 (2-carboxyethyl)-9-
 45 acridinium carboxamide
 trifluoromethanesulfonate
 [compound 61]
 MS ($M + 14$) @ 477; M^+ @ 463
 50 M_p 227°C (decomp.)

Example 7

55

Preparation of Acridinecarboxamides

An amine (110 mole percent) and triethylamine (220 mole percent) were dissolved in methylene chloride. One hundred mole percent of 9-chlorocarbonyl acridine was added dropwise as a solution in methylene chloride. The reaction was stirred under N_2 for 3 hours. The solution was filtered through silica gel and the filtrate was evaporated to leave a residue. The residue was then recrystallized from an appropriate solvent (isopropyl ether for compound 78 and ethyl ether for compound 79).

The following amides were prepared, and are described using the numerals, symbols and abbreviations which are explained in Example 1.

78. 14973-15A N-Allyl-9-acridinecarboxamide
[Allyl amine (Aldrich)]
MS M^+ @ 262

M_p 192°C

79. 14973-6A Benzyl N-(5-carboxypentyl)-9-
acridinecarboxamide
[6-Amino caproic acid (Aldrich)]
MS M^+ @ 458
 M_p 86°C

Example 8

Synthesis of Acridinium carboxamides

An ester (either compound 44 or compound 68) was added to a 1 N HCl solution and refluxed for 3-4 hours. Upon cooling, the suspension was either filtered and the product collected, or the suspension was extracted with a chloroform:isopropanol (3:2) mixture, which provided the desired product (compound 80 or 81, respectively) on evaporation. Compounds 80 and 81 are described using the numerals, symbols and abbreviations which are explained in Example 1.

80. 14379-27A 10-Methyl-N-(5-carboxypentyl)-N-
p-bromobenzenesulfonyl-9-
acridinium carboxamide
trifluoromethanesulfonate
[compound 44]
MS M^+ @ 569, 571
 M_p 148-150°C

81. 14973-51A 10-Methyl-N-(5-carboxypentyl)-N-
 p-nitrobenzenesulfonyl-9-
 acridinium carboxamide
 trifluoromethanesulfonate
 [compound 68]
 MS M⁺ @ 536

Example 9

15 Synthesis of 10-(3-sulfopropyl)-N-tosyl-N-(2-carboxyethyl)-9-acridinium carboxamide

Propane sultone (260 mole percent) was heated with t-butyl N-tosyl-N-(2-carboxyethyl)-9-acridinecarboxamide (compound 39, 13514-069) at 110-120°C for 2 hours. After cooling, the solid mass was taken up in methanol and filtered. The filtrate was evaporated to dryness and the residue triturated with benzene to remove un-quaternized material.

The crude product compound was treated with trifluoroacetic acid at 0°C then allowed to warm to 25°C over a period of 15 minutes. The residue obtained upon evaporation was purified chromatographically on preparative thick-layer chromatography plates (C-18 PLKC 18F, 20 x 20 cm, 1000M, as available from Whatman, Clifton, New Jersey), eluted with 70 parts methanol/30 parts 0.5% aqueous acetic acid, and further purified by ion exchange on Cellex-DTM resin [BioRad Laboratories, Richmond, California] using 8% formic acid to elute the product, compound 82, which is described below using the numerals, symbols and abbreviations which are explained in Example 1.

30 82. 14496-243 10-(3-sulfopropyl)-N-tosyl-N-(2-
 carboxyethyl)-9-
 acridinium carboxamide
 35 [compound 39]
 MS M⁺ @ 572

Example 10

Synthesis of 10-(3-sulfopropyl)-N-tosyl-N-(3-sulfopropyl)-9-acridinium carboxamide

45 Fifty milligrams of N-tosyl-9-acridinecarboxamide (compound 46, 14973-88A) were heated at 140-150°C under argon in a sealed tube with 500 mg of propane sultone for 3 hours. After cooling, excess propane sultone was removed by trituration with benzene (5 ml X 3). The crude product was purified by anion exchange chromatography using BioRad AG-1-X4 formate form [BioRad Laboratory, Richmond, California], eluted with a gradient of aqueous formic acid. The product, compound 83, is described below using the numerals, symbols and abbreviations explained in Example 1.

50 83. 30253-020 10-(3-Sulfopropyl)-N-tosyl-N-(3-
 sulfopropyl)-9-
 acridinium carboxamide.
 55 [compound 46]
 MS M + H @ 621.

Example 11

Evaluation of N-sulfonylacridinium carboxamide Chemiluminescence

5 Acridinium compounds to be tested for chemiluminescence were dissolved in dimethyl formamide ("DMF") and then diluted with 0.05 M sodium citrate (pH 5.0) or 0.05 M sodium phosphate (pH 7.0) buffer to give solutions of about 3×10^{-3} M. Twenty microliters of each buffered solution was diluted with 300 μ l of 0.1 N HCl and chemiluminescence was triggered with 150 μ l of 0.03% H_2O_2 in 0.2 N NaOH.

10 The light generated was recorded on a photon counter luminometer over a 10 second interval except where a longer interval is indicated in Table 1. The specific activity of each compound is provided in the form of counts/moles in Table 1.

TABLE 2

	<u>Compound No.</u>	<u>Identifying No.</u>	<u>Counts/Mole</u>
20	49	13513-246	9.4×10^{18}
	50	13513-247	9×10^{18}
	51	13513-248	1×10^{19}
25	50	13513-249	1.2×10^{19}
	53	13513-250	1×10^{19}
	54	13514-013	8.3×10^{18}
30	55	13514-014	1.25×10^{19}
	56	13514-018	1.1×10^{19}
	57	13514-021	1.5×10^{19}
35	58	13514-037	5.2×10^{18}
			(50 secs)
	59	13514-038	1.4×10^{19}
40			(20 secs)
	62	13513-211	5×10^{18}
	63	13513-212	7×10^{18}
	64	13513-215	6.1×10^{18}
45	65	13513-216	8×10^{18}
	66	13513-230	5×10^{18}

30 Example 12

Stability Test of Compound 62 (13513-211)

55 Compound 62 (2 mg) was dissolved in 1 ml of methanol. Fifty microliters of this solution were added to each of the following buffers:

- 1) 500 microliters of 0.05 M sodium phosphate, pH 5.0
- 2) 500 microliters of 0.05 M sodium phosphate, pH 5.5
- 3) 500 microliters of 0.05 M sodium phosphate, pH 6.0

4) 500 microliters of 0.05 M sodium phosphate, pH 6.5

5) 500 microliters of 0.05 M sodium phosphate, pH 7.0. Each solution was analyzed on a Perkin-Elmer Series 4 HPLC using a reverse phase column (C-18 μ Bondapak, 3.9 mm x 30 cm, available from Waters Associates, Milford, Massachusetts). The elution was done with 75% methanol and 25% 5 mM pentanesulfonic acid in 1% aqueous acetic acid at a flow rate of 1 ml/min. The effluent was monitored at 254 nm.

After 4 weeks at room temperature, the solutions at pH 5.0, pH 5.5 and pH 6.0 showed no sign of decomposition, while at pH 6.5 and at pH 7.0, 20% and 70% decomposition were seen, respectively.

10

Example 13

Comparison of Temperature and pH Stabilities of Acridinium Compounds in Buffer at pH 7.2

15 Three different acridinium compounds, compound 62, 13513-211, a compound identified by the number 13514-020 [4-(carbobenzoyloxymethyl)-phenyl-10-methyl-9-acridinium carboxylate (trifluoromethanesulfonate)] as prepared as in Weeks, et al., *Clin. Chem.*, 29, 1474-79 (1983), and compound 83, 30253-020, were compared for temperature and pH stability. The comparison was carried out in methanol or water at a concentration of 1.0 mg/ml (which is approximately equivalent to 1.6×10^{-5} M). Each of the samples was
20 diluted 1:100 in an acid solution containing one part of 0.1 N HCl plus one part phosphate-buffered saline ("PBS") pH 6.8 with 0.01% Tween 20 \oplus (available from Sigma Chemical Company, St. Louis, Missouri). The final pH of the diluent solution were about 1.5. The molarity of each of these solutions was 1.6×10^{-5} M.

Each of the solutions was scanned to record a UV-visible absorption spectrum in order to determine molar extinction coefficients and in order to detect any appreciable differences in the absorbance spectra.

25 The UV-visible absorption spectra of these acridinium compounds have the characteristics presented in Table 2.

TABLE 3

30

	Compound	Identifying	Wavelength	Observed
	No.	No.		Absorbance
35	62	13513-211	263nm	1.40
			369nm	0.286
40	83	30253-020	263.5nm	1.42
			370nm	0.304
45		13514-020	262nm	1.72
			368nm	0.334

For all three compounds, $\epsilon_{260} \approx 18,000$ and $\epsilon_{360} \approx 87,000$.

50 These spectra indicate that there is very little difference either in UV-visible absorbance or in molar extinction coefficients among these three compounds. In fact, within the limitations of experimental error, few or no spectral differences were observed.

The 1.6×10^{-5} M stock solutions of the three compounds were serially diluted 10-fold in 0.01 M sodium phosphate with 0.05% normal human serum at pH 4.8. They were also serially diluted 10-fold in PBS (pH 7.2) with 0.01% Tween 20 \oplus .

55 Because it is known that, in general, acridinium compounds are more stable at an acid pH, it was assumed that the counts obtained from the samples diluted in pH 4.8 buffer would be representative of the maximum stability with maximum chemiluminescence output. All three compounds were serially diluted 10-fold to a final concentration of 1.6×10^{-10} M. A 10 μ l aliquot of each sample was added to 90 μ l of 0.05 N

HCl. Chemiluminescence was triggered with 200 μ l of 0.03% H_2O_2 in 0.25 N NaOH and counts were monitored on a luminometer for 6 seconds with results as presented in Table 3. Results are presented in Table 3 for each of three runs.

TABLE 4

<u>Compound No.</u>	<u>Identifying No.</u>	<u>Counts/6 Seconds</u>
62	13513-211	92,669 91,241 91,995
83	30253-020	138,791 141,962 145,133
	13514-020	59,438 59,443 59,449

Within experimental error, chemiluminescent output on the luminometer did not differ among the compounds, as indicated in Table 4.

TABLE 5

Chemiluminescent Output at pH 4.8

<u>Compound No.</u>	<u>Identifying No.</u>	<u>Counts/Mole</u>
62	13513-211	5.7×10^{19}
83	30253-020	8.7×10^{19}
	13514-020	3.7×10^{19}

When 10 μ l of these same compounds were diluted to 1.6×10^{-10} M in 90 μ l PBS buffer (pH 7.2) with 0.01% Tween 20[®] and not acidified prior to running chemiluminescence output determinations as above, the results were somewhat different, especially for the acridinium carboxylate compound 13514-020, as shown in Table 5. Results are presented in Table 5 for each of three runs.

TABLE 6

Chemiluminescent Output at pH 7.2

	<u>Compound No.</u>	<u>Identifying No.</u>	<u>Counts/6 Seconds</u>
10	62	13513-211	88,633 89,135 90,394
15	83	30253-020	133,560 137,929 142,299
20		13514-020	8,185 7,274 6,363

The compound identified by the number 13514-020 produced only 4.4×10^3 counts/mole in pH 7.2 buffer, almost an order of magnitude fewer counts than it produced at pH 4.8. This may be due to pseudobase formation by a large proportion of the molecules at the more alkaline pH, the pseudobase being substantially less chemiluminescent than the corresponding positively charged acridinium compound.

The N-sulfonylacridinium carboxamide compounds showed only a very small drop in counts when incubated at pH 7.2. This suggests that they do not undergo pseudobase formation to any appreciable degree, at least at this pH.

The dilution series of all three of the acridinium compounds in pH 7.2 buffer were stored overnight at room temperature and then assayed. Both N-sulfonylacridinium carboxamide compounds showed virtually no change in chemiluminescence. The phenyl acridinium carboxylate showed a significant drop after 20 hours at room temperature.

The samples were then placed in an incubator at 45°C. Every day for the duration of the study they were removed from the incubator, cooled to room temperature, and 10 μ l aliquots diluted in 90 μ l of PBS buffer (pH 7.2) were assayed for chemiluminescence.

Neither of the N-sulfonylacridinium carboxamides showed any significant difference in chemiluminescent output when diluted either in 0.05 N HCl or in PBS at pH 7.2. However, the acridinium carboxylate 13514-020 exhibited a significantly different chemiluminescent output when diluted in 0.05 N HCl or in PBS buffer at pH 7.2. When diluted in PBS buffer (pH 7.2), the acridinium carboxylate consistently produced at least 10-fold fewer counts than when diluted in 0.05 N HCl.

The 10,N-bis-(3-sulfopropyl) acridinium carboxamide (compound 83, 30253-020) appears to be quite stable at pH 7.2 at 45°C. After 10 days under such conditions no appreciable loss of chemiluminescence was observed. Compound 13513-211 produced 10-fold fewer counts, and the acridinium carboxylate 13514-020 produced 10³ fewer counts under the same conditions.

Example 1:

Preparation of Labeled IgG

Disulfopropyl compound 83, 30253-020, was activated by treatment with phosphorous oxychloride in acetonitrile at 45°C for 12 hours under argon. The solvent and excess POCl₃ were removed in vacuo and the activated compound was used directly in the labeling reaction.

Thus, 10 mg of rabbit IgG (Sigma Chemical Company, St. Louis, Missouri) was dissolved in 0.1 M sodium phosphate buffer (2 ml, pH 7.0) containing 1% Tween 80®. One ml of this solution was mixed with about 2 mg of the bis-sulfonylchloride. The solution was agitated periodically by sonication and stirring for one hour at room temperature.

5 An aliquot (0.5 ml) of the reaction solution was chromatographed over Sephadex® G-25 (10 cm X 0.75 cm), as available from Pharmacia, Piscataway, New Jersey, and eluted with 0.1 M phosphate buffer (pH 6.5).

The labeled protein eluted as a weakly green fluorescent band. The labeled protein was further purified by HPLC using a Bio-Sil® TSK-250 column (BioRad, Richmond, California). The resulting conjugate (30253-34) contained 0.8 labels:protein, as determined from the ratio of the absorbance of 370 nm ($\epsilon = 10,000$, acridinium salt) to the absorbance 280 nm ($\epsilon = 210,000$, IgG).

Example 15

Heat Stability Studies

15 The conjugate 30253-34, as synthesized in Example 14, was serially diluted 10-fold in three buffers (0.1 M sodium phosphate, 0.01% Tween 20®, pH 6.3; 0.01 M sodium phosphate, 0.15 M NaCl, 0.01% Tween 20®, pH 6.8; and 0.01 M sodium phosphate, 0.15 M NaCl, 0.01% Tween 20®, pH 7.2) to a concentration of 2×10^{-7} M IgG and 1.6×10^{-7} M acridinium. A dilution series was prepared and initial counts were recorded by taking 10 μ l of the sample, diluting with 90 μ l of PBS buffer at pH 6.3, pH 6.8, or pH 7.2, and then triggering chemiluminescence with 200 μ l of 0.03% H₂O₂ in 0.25 N NaOH. A 100 MI sample of PBS buffer was used as a control for each series.

25 Counts shown in Table 6 are averages of results for duplicate samples assayed on the day on which the dilution series was prepared. The concentration shown in Table 5 is the concentration of the sample prior to dilution. The amount of parentheses for each entry in Table 5 is the amount of conjugate present in the sample.

TABLE 7

	<u>Concentration (Amount)</u>	<u>Counts/6 Seconds</u>
5	<u>pH 6.3</u>	
	buffer (0 moles)	253
10	2 X 10 ⁻¹⁰ M (2 X 10 ⁻¹⁴ moles)	216,054
	1 X 10 ⁻¹⁰ M (1 X 10 ⁻¹⁴ moles)	100,842
	5 X 10 ⁻¹¹ M (5 X 10 ⁻¹⁵ moles)	48,704
15	2.5 X 10 ⁻¹¹ M (2.5 X 10 ⁻¹⁵ moles)	23,771
	1.25 X 10 ⁻¹¹ M (1.25 X 10 ⁻¹⁵ moles)	11,475
	6 X 10 ⁻¹² M (6 X 10 ⁻¹⁶ moles)	5,866
20	<u>pH 6.8</u>	
	buffer (0 moles)	233
	2 X 10 ⁻¹⁰ M (2 X 10 ⁻¹⁴ moles)	295,608
25	1 X 10 ⁻¹⁰ M (1 X 10 ⁻¹⁴ moles)	149,725
	5 X 10 ⁻¹¹ M (5 X 10 ⁻¹⁵ moles)	76,820
	2.5 X 10 ⁻¹¹ M (2.5 X 10 ⁻¹⁵ moles)	38,801
30	1.25 X 10 ⁻¹¹ M (1.25 X 10 ⁻¹⁵ moles)	18,408
	6 X 10 ⁻¹² M (6 X 10 ⁻¹⁶ moles)	9,398
35	<u>pH 7.2</u>	
	buffer (0 moles)	726
	2 X 10 ⁻¹⁰ M (2 X 10 ⁻¹⁴ moles)	309,445
	1 X 10 ⁻¹⁰ M (1 X 10 ⁻¹⁴ moles)	156,311
40	5 X 10 ⁻¹¹ M (5 X 10 ⁻¹⁵ moles)	77,238
	2.5 X 10 ⁻¹¹ M (2.5 X 10 ⁻¹⁵ moles)	39,879
	1.25 X 10 ⁻¹¹ M (1.25 X 10 ⁻¹⁵ moles)	19,925
45	6 X 10 ⁻¹² M (6 X 10 ⁻¹⁶ moles)	10,526

Each dilution series was placed in a warm air incubator at 45°C after an initial reading was taken. A duplicate reading was made on each sample daily and then the readings were averaged.

When the conjugate was stored at pH 6.8 and at 45°C, there was no loss in chemiluminescent activity of the label over a 15 day period of observation, at any dilution. Essentially the same results were observed when the conjugate was stored in PBS buffer at pH 7.2.

Example 16

35 Comparison of CLIA vs. RIA

A. Preparation of Acridinium-Labeled Anti-HsAg Conjugate.

Compound 75 (13514-081, Example 6) (12.5 μ mol) was dissolved in 200 μ l of DMF, was treated with NHS (dissolved in 50 μ l of DMF) and dicyclohexylcarbodiimide (dissolved in 50 μ l of DMF) ("DCC"); and stirred for 12 hours at room temperature. The solution of the activated ester was mixed with mouse monoclonal anti-HBsAg in 0.1 M sodium phosphate buffer (pH 6.3) in a molar ratio of 100:1 at 4°C for 12 hours.

The conjugate was then dialysed against PBS buffer, pH 6.3, until the absorbance of the dialysate indicated no free label. A UV spectral analysis indicated between 2 to 6 labels/antibody (as determined from a ratio of absorbances as in Example 14).

B. Assay for HBsAg.

Either type A_d or type A_y HBsAg (200 μ l) was diluted in calf serum and was reacted with an Auszyme™ (Abbott Laboratories, Abbott Park, Illinois) monoclonal antibody bead and 2×10^5 of counts of ¹²⁵I-labeled mouse monoclonal anti-HBsAg antibody (40 μ l, in the RIA) or an acridinium-labeled mouse monoclonal anti-HBsAg antibody (40 μ l, in the CLIA) in PBS containing 50% calf serum, 10% human serum, 0.05% Tween 20 and 5 mM EDTA (pH 6.3), for three hours at 40°C. The beads were then washed 6 times in water and counted for their activities. Calf serum was used as a negative control.

In the CLIA, a polystyrene bead with conjugate bound adsorbed thereto was mixed with 250 μ l phosphate, 0.5 mM, pH 5.3, in a glass vial suitable for use in a luminometer. While the sample was in the measuring position, 0.2 ml of 0.03% H₂O₂ in 0.25 N NaOH was then injected into the glass vial. The light emitted was measured in the luminometer. Reading began 0.012 seconds before initiation of the chemical reaction and continued for 6 seconds.

The results are presented in Table 7.

TABLE 8

Concentration (ng/ml)	CLIA		RIA	
	A _d	A _y	A _d	A _y
1.0	2214	3144	371	400
0.5	1256	2494	236	408
0.25	701	921	221	248
0.125	521	592	173	179
Calf Serum	151	179		
Cut-off	327	376		

Under the stated conditions, the sensitivity for the CLIA was less than 0.125 ng/ml for both the A_d and A_y types of HBsAg. For the RIA the sensitivity was 1.0 ng/ml for both the A_d and A_y types. The cut-off count was 2.1 times that of the negative control.

Table 8 clearly shows that chemiluminescent immunoassays according to the present invention are more sensitive than comparable radioimmunoassays.

Example 17

A comparison of CLIA and EIA

A. Preparation of labeled anti-hTSH (30234-207).

Compound 75 (13514-081, Example 6) (2 mg, 4.3 μ moles) in 200 ml of acetonitrile was treated with 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide hydrochloride (Sigma, St. Louis, Missouri) (10 μ moles) in 100 μ l of acetonitrile and N-hydroxysuccinimide (4.0 μ moles) in 100 μ l of acetonitrile for 12 hours at 25°C in the dark.

The active ester was mixed with anti-hTSH in PBS buffer containing 0.5% 3-[(3-cholamidopropyl) dimethylammonio]-1-propane-sulfonate ("CHAPS") at pH 6.5 in a ratio of 50:1 (antibody:active ester). After coupling for 3 hours at 25°C, the labeled antibody was dialysed against PBS buffer containing 0.5% CHAPS at pH 6.5 until no free label was present in the dialysate by U.V.

Based on the U.V. spectra, the conjugate had an average of 10 labels per antibody.

B. Assay for hTSH.

CLIA and EIA were compared using the Abbott hTSH-EIA Kit (Abbott Laboratories, Abbott Park, Illinois) with the exception that for the CLIA, the anti-hTSH acridinium conjugate was used in place of the kit anti-hTSH-HRPO conjugate. Thus, a standard curve was generated by incubating the kit standards with the kit beads at 37°C for 1 hour, then washing three times. For the CLIA, the conjugate prepared above was diluted 1:5000 with PBS buffer containing 50% calf serum, 1% normal mouse serum, 0.05% Tween® 20 and 2 mM EDTA at pH 6.3. One hundred microliters of this solution was incubated with the beads for 1 hour at 37°C, then washed four times.

The beads were transferred one by one to the reaction vial of a luminometer containing 400 μ l of water and reacted with 200 μ l of 0.03% H₂O₂ in 0.2 N NaOH. Photon counts were recorded for 6 seconds.

The EIA was carried out according to the instructions in the kit insert on a Quantum II® spectro photometer (Abbott Laboratories, Abbott Park, Illinois)

The results are shown in Table 9.

TABLE 9

Concentration (μ Iu/ml)	CLIA (counts)	EIA (A ₄₉₂)
0	533 (SD 35.4)	0.012
1	5064	0.062
4	14476	0.176
10	32092	0.397
25	66072	0.828
60	110,984	1.602

Under these conditions the sensitivity of the CLIA was 0.016 μ IU/ml (0 standard + 2 SD) while the EIA had a sensitivity of 0.05 μ IU/ml.

Example 18

Preparation of 5-methyl-6-[N-tosyl-N-(2-carboxyethyl)]-phenanthridiniumcarboxamide

Phenanthridine-6-carboxylic acid (400 mg, 1.8 mmoles) [prepared by the method of Wittig et al., *Justus Liebig's Ann.*, 577, 1 (1952)], was suspended in methylene chloride (20 ml, distilled from P₂O₅) and cooled to 0°C under nitrogen. Oxalyl chloride (320 μ l, 3.6 mmoles) (Aldrich Chemical Co., Milwaukee, Wisconsin) was added, followed by DMF (5 μ l). As the reaction mixture was stirred for one hour at 0°C and for 30

minutes at 25°C, all the carboxylic acid dissolved. The solution was evaporated to dryness to give the acid chloride which was used without further purification.

Methyl N-tosyl-β-alanine was prepared from methyl-β-alanine (Aldrich Chemical Company, Milwaukee, Wisconsin) and tosyl chloride (Aldrich Chemical Company, Milwaukee, Wisconsin) according to the procedure of Example 1. Potassium t-butoxide (600 mg, 5.4 mmoles, freshly sublimed) was added to a solution of 1.3g (5.4 mmoles) of methyl N-tosyl-β-alanine in 50 ml of THF. After stirring for 15 minutes and at room temperature and under N₂, the suspension was evaporated to dryness. The potassium salt of methyl N-tosyl-β-alanine, was resuspended in 20 ml of THF, mixed with the acid chloride (in 20 ml of THF), and stirred for 12 hours.

The resulting suspension was poured into 100 ml of ethylacetate, washed with 50 ml of water and washed twice with 25 ml of brine. After drying over MgSO₄ and evaporating to dryness, the residue was chromatographed on a Chromatatron™ chromatograph (available from Harrison Research, Palo Alto, California) using a 4 mm silica rotor and employing a 25/75 ethylacetate/hexane gradient. The product (R_f 0.2) was collected, then recrystallized from benzene/hexane (i.e., the product was dissolved in benzene, and hexane was added until cloudy) to give 130 mg of methyl 6-[N-tosyl-N-(2-carboxyethyl)]-phenanthridinecarboxamide, Compound 84, 13514-225.

Compound 84, 13514-225, was methylated according to the procedure in Example 3 to give methyl 5-methyl-6-[N-tosyl-N-(2-carboxyethyl)]-phenanthridiniumcarboxamide, compound 85, 13514-227. Compound 85 was hydrolyzed according to the procedure in Example 8 to provide 5-methyl-6-[N-tosyl-N-(2-carboxyethyl)]-phenanthridiniumcarboxamide, compound 86, 13514-228.

Compounds 84, 85 and 86 are described using the numerals, symbols and abbreviations as explained in Example 1.

84. 13514-225 Methyl 6-[N-tosyl-
N-(2-carboxyethyl)]-
phenanthridinecarboxylate
MS M + H @ 463

85. 13514-227 Methyl 5-methyl-
6-[N-tosyl-N-
(2-carboxyethyl)]-
phenanthridiniumcarboxamide
MS M⁺ @ 477
Mp 136°C


86. 13514-228 5-Methyl-6-[N-tosyl-
N-(2-carboxyethyl)]-
phenanthridiniumcarboxamide
MS M⁺ @ 463

Although the present invention has been described in terms of preferred embodiments, it is understood that modifications and improvements will occur to those skilled in the art.

For example, in light of the results presented herein, it is expected that additional compounds which are useful according to the present invention may be identified by the formula

5

10




wherein a, b, c, d, a', b', c', d' independently may be hydrogen, alkyl, aryl, amino, substituted amino,
 15 carboxy-alkyl, sulfoalkyl, alkoxy, aryloxy, sulfo, thio alkoxy, thioaryloxy aminoalkyl, protected aminoalkyl,
 hydroxyalkyl, protected hydroxyalkyl, haloalkyl, or any adjacent of these positions may be linked so as to
 form aromatic rings fused to the acridine nucleus.

In addition, Sheehan et al., U.S. Patent No. 3,539,574 described chemiluminescent acridinium com-
 pounds which are also expected to be useful according to the present invention. Other isomeric acridinecar-
 20 boxylic acids, quinoline carboxylic acids, isoquinoline carboxylic acid, other activated acridine amides, and
 other activated acridine esters are expected to be useful according to the present invention. Such
 compounds include, without limitation: hydroxamates identified by the formula

25

30

35




enamides identified by the formula

40

45


50



arylamides identified by the formula

55


5

Y⁻

10

wherein X and Y⁺ are electron withdrawing groups; N-heterocycles identified by the formula

15


Y⁻

20

25

30 wherein X and U⁺ may independently be O, S, P, N, or C; activated esters such as thioesters identified by the formula


35



40

or such as thioesters identified by the formula

45


Y⁻

50

55

acridine acids identified by the formula

5




Y -

10

or by the formula


15



20

quinoline acids identified by the formula


25



30

or by the formula


35



40

or isoquinoline acids identified by the formula


45



50

55

or by the formula



It is understood that those skilled in the art will be enabled by the above specification to incorporate reactive functional groups for attaching the label to an analyte into compounds according to the present invention.


It is also contemplated that compounds according to the present invention will be: used in labeling DNA probes; incorporated into an enzyme substrate wherein the product of the enzymatic reaction is the chemiluminescent compound; and incorporated into systems which involve energy transfer or fluorescent quenching.

Compounds according to the present invention may also be: incorporated into a system which employs the compound as a labeling reagent in a post-column HPLC detection system; used to measure H_2O_2 concentration; and used as a source of intense pulsed light.

Therefore, it is intended that the present invention include all such variations and improvements as come within the scope of the invention as claimed.

Claims

1. a chemiluminescent compound identified by the formula



and isomers thereof wherein R, R', R'', X', X2, and X3 are substituents which do not interfere with effective chemiluminescence with the proviso that R-X3, R'-X2 and R''-X' may independently be hydrogen.

2. The chemiluminescent compound as recited in claim 1:

wherein R, R', and R'' are spacer arms;

wherein X', X2 and X3 are independently members of the group consisting of hydrogen, carboxy, carboalkoxyl, carboxamido, carboaryloxy, cyano, carboximido, isocyanato, isothiocyanato, sulfo, sulfonyl halide, carbonyl halide, N-succinimidylcarboxy and N-maleimide groups; and

wherein Y is an appropriate counter ion.

3. The chemiluminescent compound as recited in claim 2 wherein Y is a counter ion selected from the group consisting of sulfate, alkylsulfate, halosulfate, haloborate, haloacetate, halophosphate, phosphate and halide.

4. The chemiluminescent compound as recited in claim 1 wherein R, R', and R'' independently comprise a member selected from the group consisting of alkyl, alkylene, aryl, substituted alkyl, substituted alkylene and substituted aryl groups such that:

one or more hydrogens of said member is replaced by an alkyl, aryl, alkylene, substituted alkyl, substituted alkylene, substituted, aryl, alkoxy, aryloxy, halo, amino protected amino, substituted amino hydroxy, protected hydroxy, oxo, thio, imino, mercapto or substituted mercapto group;

or such that one or more carbon atoms of the member is replaced by a heteroatom.

5. The chemiluminescent compound as recited in claim 4 wherein said heteroatom is selected from the group consisting of nitrogen, phosphorus, sulfur and oxygen.


6. The chemiluminescent compound as recited in claim 2 wherein R, R', and R'' independently are spacer arms of the formula

5 $-(CH_2)_n-$

where n = 0 - 50.

7. The chemiluminescent compound as recited in claim 1 wherein R'' is $-CH_2-$, X' is $-H$, and R'-X² is identified by the formula

10



15 8. The chemiluminescent compound as recited in claim 7 wherein said compound is identified by the formula 10-methyl-N-[2-carboxyethyl]-N-tosyl-9-acridinium carboxamide.


9. The chemiluminescent compound as recited in claim 7 wherein said compound is identified by the formula 10-methyl-N-(4-carboxybutyl)-N-tosyl-9-acridinium carboxamide.

10. The chemiluminescent compound as recited in claim 7 wherein said compound is identified by the formula 10-methyl-N-(5-carboxypentyl)-N-tosyl-9-acridinium carboxamide.

20

11. The chemiluminescent compound as recited in claim 1 wherein formula R'' is $-(CH_2)_7-$, X' is $-SO_2-$, and R'-X² is identified by the formula

25




12. The chemiluminescent compound as recited in claim 11 wherein said compound is identified by the formula 10-(3-sulfopropyl)-N-(2-carboxyethyl)-N-tosyl-9-acridinium carboxamide.

13. The chemiluminescent compound as recited in claim 11 wherein said compound is identified by the formula 10-(3-sulfopropyl)-N-(3-sulfopropyl)-N-tosyl-9-acridinium carboxamide.


14. The chemiluminescent compound as recited in claim 7 wherein R'-X² is identified by the formula.

35



and wherein R-X³ is identified by the formula

40



45

15. The chemiluminescent compound as recited in claim 7 wherein said compound is selected from 10-methyl-N-phenyl-N-tosyl-9-acridinium carboxamide trifluoromethanesulfonate, 10-methyl-N-phenyl-N-(p-bromobenzenesulfonyl)-9-acridinium carboxamide trifluoromethanesulfonate, 10-methyl-N-phenyl-N-(p-nitrobenzenesulfonyl)-9-acridinium carboxamide trifluoromethanesulfonate, 10-methyl-N-phenyl-N-(o-nitrobenzenesulfonyl)-9-acridinium carboxamide trifluoromethanesulfonate, or 10-methyl-N-phenyl-N-trifluoromethanesulfonyl-9-acridinium carboxamide trifluoromethanesulfonate.


55

16. The chemiluminescent compound as recited in claim 7 wherein said compound is identified by the formula 10-methyl-N-isopropyl-N-tosyl-9-acridinium carboxamide trifluoromethanesulfonate, 10-methyl-N-isopropyl-N-(p-bromobenzenesulfonyl)-9-acridinium carboxamide trifluoromethanesulfonate, 10-methyl-N-isopropyl-N-(o-nitrobenzenesulfonyl)-9-acridinium carboxamide trifluoromethanesulfonate, or 10-methyl-N-isopropyl-N-trifluoromethanesulfonyl-9-acridinium carboxamide trifluoromethanesulfonate.


17. The chemiluminescent compound as recited in claim 7 wherein said compound is identified by the formula 10-methyl-N-butyl-N-(2,4,6-trimethylbenzenesulfonyl)-9-acridinium carboxamide trifluoromethanesulfonate, 10-methyl-N-butyl-N-(2,4,6-tri-isopropyl-benzenesulfonyl)-9-acridinium carboxamide trifluoromethanesulfonate, 10-methyl-N-phenyl-N-tosyl-9-acridinium carboxamide trifluoromethanesulfonate, 10-methyl-N-butyl-N-(p-bromobenzenesulfonyl)-9-acridinium carboxamide trifluoromethanesulfonate, 10-methyl-N-butyl-N-(o-nitrophenylsulfonyl)-9-acridinium carboxamide trifluoromethanesulfonate, 10-methyl-N-butyl-N-(p-nitrobenzenesulfonyl)-9-acridinium carboxamide trifluoromethanesulfonate, 10-methyl-N-butyl-N-(2,4-dinitrobenzenesulfonyl)-9-acridinium carboxamide trifluoromethanesulfonate, or 10-methyl-N-allyl-N-tosyl-9-acridinium carboxamide trifluoromethanesulfonate.

18. The chemiluminescent compound as recited in claim 1 wherein said compound is identified by the formula methyl 6-[N-tosyl-N-(2-carboxyethyl)]-phenanthridinecarboxamide, methyl 5-methyl-6-[N-tosyl-N-(2-carboxyethyl)]-phenanthridiniumcarboxamide, or 5-methyl-6-[N-tosyl-N-(2-carboxyethyl)]-phenanthridiniumcarboxamide.

19. A method for preparation of a chemiluminescent compound comprising the steps of:
 contacting an amine identified by the formula $R-X^3NH_2$
 with a sulfonylhalide identified by the formula $W-SO_2-R'-X^2$
 in an inert solvent in the presence of base to form a sulfonamide anion and metal ion identified by the formulas



a) acylating with an activated 0-acridinecarboxylic acid identified by the formula



wherein R, R' and R'' are independently selected from the group consisting of: alkyl, aryl, alkylene, substituted alkyl, substituted alkylene and substituted aryl, alkoxy, aryloxy, halo, amino, protected, amino, substituted amino hydroxy, protected hydroxy, oxo, thio, imino, mercapto groups; and alkyl, aryl, alkylene substituted alkyl, substituted alkylene and substituted aryl groups comprising a heteroatom selected from the group consisting of nitrogen, phosphorus, sulfur and oxygen with the proviso that R-X³, R-X², and R''-X' may independently be hydrogen;

wherein X¹, X², and X³ are independently members of the group consisting of hydrogen, carboxy, carboalkoxyl, carboxamido, carboaryloxy, cyano, carboximido, isocyanato, isothiocyanato, sulfo, sulfonyl halide, carbonyl halide, N-carboxysuccinimide and N-maleimide groups;


wherein Y is an appropriate counter ion;

wherein, W is selected from the group consisting of chloro and fluoro groups; and

wherein M is selected from the group consisting of Li, Na and K; and

wherein Z is selected from the group consisting of halo, imidazo, N-hydroxysuccinimidyl and azido groups; or

b) acylating with an activated phenanthridine-6-carboxylic acid identified by the formula



wherein R, R' and R'' are independently selected from the group consisting of: alkyl, aryl, alkylene, substituted alkyl, substituted alkylene and substituted aryl, alkoxy, aryloxy, halo, amino, protected amino, substituted amino hydroxy, protected hydroxy, oxo, thio, imino, mercapto groups; and alkyl, aryl, alkylene substituted alkyl, substituted alkylene and substituted aryl groups comprising a heteroatom selected from the group consisting of nitrogen, phosphorus, sulfur and oxygen with the proviso that R-X³, R'-X and R''-X may independently be hydrogen;

wherein X¹, X² and X³ are independently members of the group consisting of hydrogen, carboxy, carboalkoxyl, carboxamido, carboaryloxy, cyano, carboximido, isocyanato, isothiocyanato, sulfo, sulfonyl halide, carbonyl halide, N-carboxysuccinimide and N-maleimide groups;

wherein Y is an appropriate counter ion;


wherein W is selected from the group consisting of chloro and fluoro groups; and

wherein M is selected from the group consisting of Li, Na and K, and




wherein Z is selected from the group consisting of halo, imidazolo, N-hydroxysuccinimidyl and azido groups.

20. A conjugate formed by an antibody or antigen conjugated to a chemiluminescent compound as recited in claim 1.

21. A method for performing a chemiluminescent immunoassay to test for the presence of an antigen or antibody to an antigen as recited in claim 20 comprising the step of exposing a sample to a conjugate as recited in claim 20.



HCl ,

 $\text{Y}-\text{R}''-\text{X}^1$  Y^-